
 NOR-CAL EMS	05-0810	Pediatric Non-Traumatic Shock
Nor-Cal EMS Policy & Procedure Manual		Name of the Section
Effective Date: 02/01/2021		Next Revision: 02/01/2023
Approval: Jeffrey Kepple MD – MEDICAL DIRECTOR		SIGNATURE ON FILE

Authority

Health and Safety Code Division 2.5, California Code of Regulations, Title 22, Division 9.



PUSH DOSE EPINEPHRINE MIXING INSTRUCTIONS (1mcg/ml):

1. Prepare a 10ml syringe with 9ml of saline.
2. Draw 1ml of Epinephrine 1:10,000
3. Label syringe Epi 10mcg/ml

OR

1. Take Epinephrine 1:10,000 concentration (1 mg/10 ml) and waste 9 ml of Epinephrine
2. In same syringe draw 9 ml of saline from the patient's IV bag & shake well
3. Label syringe Epi 10mcg/ml

Pediatric Blood Pressures

Potentially Hypotensive if Systolic BP less than:

Infant	less than 60
Toddler (1-3 yrs)	less than 70
Preschool (3-6 yrs)	less than 75
School age (6-12 yrs)	less than 80
Adolescent (12-18 yrs)	less than 90

Use formula: (age x 2) + 65 to determine if patient is potentially hypotensive
Formula for normal systolic patient: (age x 2) + 90